

Ecological Pyramids

An important feature of energy flow is that most of the energy going from one trophic level to the next in a food chain or food web dissipates into the environment as a result of the second law of thermodynamics. **Ecological pyramids** often graphically represent the relative energy values of each trophic level. There are three main types of pyramids—a pyramid of numbers, a pyramid of biomass, and a pyramid of energy.

A **pyramid of numbers** shows the number of organisms at each trophic level in a given ecosystem, with greater numbers illustrated by a larger area for that section of the pyramid (Figure 3.12). In most pyramids of numbers, the organisms at the base of the food chain are the most abundant, and fewer organisms occupy each successive trophic level. In African grasslands the number of herbivores, such as zebras and wildebeests, is far greater than the number of carnivores, such as lions. Inverted pyramids of numbers, in which higher trophic levels have *more* organisms than lower trophic levels, are often observed among decomposers, parasites, tree-dwelling herbivorous insects, and similar organisms. One tree may provide food for thousands of leaf-eating insects, for example. Pyramids of numbers are of limited usefulness because they do not indicate the biomass of the organisms at each level, and they do not indicate the amount of energy transferred from one level to another.

A **pyramid of biomass** illustrates the total biomass at each successive trophic level. **Biomass** is a quantitative estimate of the total mass, or amount, of living material; it indicates the amount of fixed energy at a particular time. Biomass units of measure vary: Biomass is represented as total volume, as dry weight, or as live weight. Typically, pyramids of biomass illustrate a progressive reduction of biomass in succeeding trophic levels (Figure 3.13). For example, if one assumes there is about a 90% reduction of biomass for each trophic level, 10,000 kg of grass should support 1000 kg of grasshoppers, which in turn support 100 kg of toads. The 90% reduction in biomass is used for illustrative purposes only; actual field numbers for biomass reduction in nature vary widely. By this logic, however, the biomass of toad eaters such as snakes could be, at most, only about 10 kg. From this brief exercise, it is apparent that although carnivores do not eat vegetation, a great deal of vegetation is required to support them.

A **pyramid of energy** illustrates the energy content, often expressed as kilocalories per square meter per year, of the biomass of each trophic level (Figure 3.14). These pyramids always have large energy bases and get progressively smaller through succeeding trophic levels. Energy pyramids show that most energy dissipates into the environment when going from one trophic level to the next: Less energy reaches each successive trophic level from the level beneath it because organisms at the lower level use some energy to perform work, and some is lost. (Remember, because of the second law of thermodynamics, no biological process is ever 100% efficient.) Energy pyramids explain why there are so few trophic levels: Food webs are short because of the dramatic reduction in energy content at each trophic level. (See “You Can Make a Difference: Vegetarian Diets” in Chapter 19 for a discussion of how the eating habits of humans relate to food chains and trophic levels.)




Figure 3.12 Pyramid of numbers. This pyramid is for a hypothetical area of temperate grassland; in this example, 10,000 grass plants support 10 mice, which support one bird of prey. Based on the number of organisms found at each trophic level, a pyramid of numbers is not as useful as other ecological pyramids. It provides no information about biomass differences or energy relationships between one trophic level and the next. (Note that decomposers are not shown.)




Figure 3.13 Pyramid of biomass. This pyramid is for a hypothetical area of temperate grassland. Based on the biomass at each trophic level, pyramids of biomass generally have a pyramid shape with a large base and progressively smaller areas for each succeeding trophic level. (Note that decomposers are not shown.)